

Fuzzy Unification and Generalization of First-Order Terms over Similar Signatures

A Constraint-Based Approach

Hassan Aït-Kaci

Gabriella Pasi

27th LOPSTR

**Namur, Belgium
October 10–12, 2017**

This presentation's objective

- ▶ **Reformulate and extend** general results on (crisp & fuzzy) *FOT* **unification and generalization** (“*anti-unification*”) seen as **lattice operations using** (crisp & fuzzy) **constraints**
- ▶ **Give declarative rulesets** for operational constraint-driven deductive and inductive **fuzzy inference over *FOTs*** when **some signature symbols may be similar**

OK... *And why is this interesting?...*

- ▶ This provides a **formally clean** and **practically efficient** way to enable ***approximate reasoning*** (**deduction** and **learning**) ***with a very popular data structure*** used in logic-based data and knowledge processing systems

Some quick but important remarks about this presentation

We apologize in advance for the “*symbol soup*” in this talk ...

... but please do bear with us, as **this presentation is:**

- ▶ **only meant to give you an idea...** of what's in the **paper** with more examples and all proofs available **here**
- ▶ **necessary...** since we purport to be formal
- ▶ **not that complicated...** at least not for this audience — *we assume familiarity with Prolog's basic data structure and Fuzzy Logic notions*
- ▶ **really always the same...** once we get the basic gist

Presentation outline

- ▶ **First-Order Terms** — syntax of $FOTs$
- ▶ **Subsumption** — pre-order relation on $FOTs$
- ▶ **Unification** — glb operation on $FOTs$
- ▶ **Generalization** — lub operation on $FOTs$
- ▶ **Weak unification** — fuzzy glb of aligned $FOTs$
- ▶ **Weak generalization** — fuzzy lub of aligned $FOTs$
- ▶ **Full fuzzy unification** — fuzzy glb of misaligned $FOTs$
- ▶ **Full fuzzy generalization** — fuzzy lub of misaligned $FOTs$
- ▶ **Conclusion** — recapitulation and future work

The lattice of \mathcal{FOTs}



data structures that can be approximated



\mathcal{FOTs} on a signature of data constructors $\Sigma \stackrel{\text{def}}{=} \bigcup_{n \geq 0} \Sigma_n$

$$\mathcal{T}_{\Sigma, \mathcal{V}} \stackrel{\text{def}}{=} \mathcal{V}$$

$$\cup \{ f(t_1, \dots, t_n) \mid f \in \Sigma_n, n \geq 0,$$

$$t_i \in \mathcal{T}_{\Sigma, \mathcal{V}}, 1 \leq i \leq n \}$$

\mathcal{FOT} subsumption pre-order relation

$$t_1 \preceq t_2$$


iff

$$\exists \sigma : \mathcal{V} \rightarrow \mathcal{T}_{\Sigma, \mathcal{V}}$$

s.t.

$$t_1 = t_2 \sigma$$

\mathcal{FOT} subsumption lattice operations



Declarative lattice operations on \mathcal{FOT} s...



using constraints




- ▶ **1930** – Jacques Herbrand gives normalization rules for sets of term equalities in his PhD thesis (*Chap. 5, Sec. 2.4, pp. 95 – 96*) but does not call this “*unification*”
- ▶ **1960** – Dag Prawitz expresses this as reduction rules as part of proof normalization procedure for Natural Deduction in F.O. Logic (Gentzen, 1934)
- ▶ **1965** – J. Alan Robinson gives a procedural algorithm and uses it to lift the resolution principle from Propositional Logic to F.O. Logic — calling it “*unification*”
- ▶ **1967** – Jean van Heijenoort translates Chap. 5 of Herbrand’s thesis into English
- ▶ **1971** – Warren Goldfarb translates Herbrand’s full thesis into English

- ▶ **1976** – **Gérard Huet** dates the first \mathcal{FOT} unification algorithm to initial equation normalization in Herbrand's 1930 PhD thesis (*also in Chap. 5 in Huet's thesis!*)
- ▶ **1982** – **Alberto Martelli & Ugo Montanari** give unification rules (with no mention of Herbrand's thesis, although Huet's thesis is cited)

Interestingly, Martelli & Montanari use a preprocessing method that uses generalization implicitly (to compute “*common parts*” in preprocessing equations into congruence classes of equations called “*multi-equations*”) — **but do not point out that it is dual to unification**

\mathcal{FOT} unification as a constraint



Declarative unification rule

A unification rule **rewrites** a **prior set of equations** E into a **posterior set of equations** E' whenever an **optional meta-condition** holds:

RULE NAME:

Prior set of equations E [*Optional meta-condition*]

Posterior set of equations E'

TERM DECOMPOSITION:

$$\frac{E \cup \{ f(s_1, \dots, s_n) \doteq f(t_1, \dots, t_n) \}}{E \cup \{ s_1 \doteq t_1, \dots, s_n \doteq t_n \}} \quad [n \geq 0]$$

VARIABLE ELIMINATION:

$$\frac{E \cup \{ X \doteq t \}}{E[X \leftarrow t] \cup \{ X \doteq t \}} \quad \left[\begin{array}{l} X \notin \mathbf{Var}(t) \\ X \text{ occurs in } E \end{array} \right]$$

EQUATION ORIENTATION:

$$\frac{E \cup \{ t \doteq X \}}{E \cup \{ X \doteq t \}} \quad [t \notin \mathcal{V}]$$

VARIABLE ERASURE:

$$\frac{E \cup \{ X \doteq X \}}{E}$$

Moving on to...



declarative constraint-based generalization



Generalization

a bit of history

- ▶ The lattice-theoretic properties of $FOTs$ as data structures pre-ordered by subsumption were exposed independently and simultaneously by **Reynolds** and **Plotkin** in **1970**
- ▶ Both gave a formal definition of FOT generalization and each proved correct a *procedural* specification for computing it
- ▶ *However*, ... so far, a **declarative** formal specification was lacking — **which we provide here**
- ▶ **Why should we care?...** Well, because:
 - **syntax-driven rules give an operational semantics as constraint solving needing no control specification** (use any rule that applies in any order)
 - **each rule's correctness is independent of that of the others** (they share no global context)
 - **eases the formal specification of more expressive approximation over the same data structure** (such as **fuzzy constraints** on $FOTs$)

\mathcal{FOT} generalization judgment

Statement of the form:

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \vdash \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} t \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$$

where (for $i = 1, 2$):

- $t \in \mathcal{T}$ and $t_i \in \mathcal{T}$ are \mathcal{FOT} s
- $\sigma_i : \mathcal{V} \rightarrow \mathcal{T}$ and $\theta_i : \mathcal{V} \rightarrow \mathcal{T}$ are substitutions

\mathcal{FOT} generalization judgment validity

A generalization judgment:


$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \vdash \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} t \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$$

is deemed **valid** whenever:

$$t_i \sigma_i = t \theta_i$$

with $\theta_i \preceq \sigma_i$ (i.e., $\exists \delta_i$ s.t. $\theta_i = \delta_i \sigma_i$) for $i = 1, 2$

\mathcal{FOT} generalization judgment validity as a constraint



Declarative generalization axiom

Statement of the form:

AXIOM NAME:

[Optional meta-condition]

Judgment J

which reads:

“whenever the optional meta-condition holds, judgement J is always valid”

\mathcal{FOT} generalization axioms

EQUAL VARIABLES :

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \vdash \begin{pmatrix} X \\ X \end{pmatrix} X \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}$$

VARIABLE-TERM :

$[t_1 \in \mathcal{V} \text{ or } t_2 \in \mathcal{V}; t_1 \neq t_2; X \text{ is new}]$

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \vdash \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} X \begin{pmatrix} \sigma_1 \{ t_1 / X \} \\ \sigma_2 \{ t_2 / X \} \end{pmatrix}$$

UNEQUAL FUNCTORS :

$[m \geq 0, n \geq 0; m \neq n \text{ or } f \neq g; X \text{ is new}]$

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \vdash \begin{pmatrix} f(s_1, \dots, s_m) \\ g(t_1, \dots, t_n) \end{pmatrix} X \begin{pmatrix} \sigma_1 \{ f(s_1, \dots, s_m) / X \} \\ \sigma_2 \{ g(t_1, \dots, t_n) / X \} \end{pmatrix}$$

Declarative generalization inference rule

Conditional Horn rule of generalization judgments of the form:

RULE NAME:

[Optional Meta-Condition]

Prior Judgment J_1 ... Prior Judgment J_n

Posterior Judgment J

(for $n \geq 0$) — which reads:

“whenever the *optional meta-condition* holds, if all the n *prior judgements J_n* are valid, then the *posterior judgement J* is also valid”

Declarative FOT generalization rule for equal functors

EQUAL FUNCTORS :

$[n \geq 0]$

$$\frac{\begin{array}{c} \left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right) \vdash \left(\begin{array}{c} s'_1 \\ t'_1 \end{array} \right) u_1 \left(\begin{array}{c} \sigma_1^1 \\ \sigma_2^1 \end{array} \right) \quad \dots \quad \left(\begin{array}{c} \sigma_1^{n-1} \\ \sigma_2^{n-1} \end{array} \right) \vdash \left(\begin{array}{c} s'_n \\ t'_n \end{array} \right) u_n \left(\begin{array}{c} \sigma_1^n \\ \sigma_2^n \end{array} \right) \end{array}}{\left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right) \vdash \left(\begin{array}{c} f(s_1, \dots, s_n) \\ f(t_1, \dots, t_n) \end{array} \right) f(u_1, \dots, u_n) \left(\begin{array}{c} \sigma_1^n \\ \sigma_2^n \end{array} \right)}$$

where $\left(\begin{array}{c} s'_i \\ t'_i \end{array} \right) \stackrel{\text{def}}{=} \left(\begin{array}{c} s_i \\ t_i \end{array} \right) \uparrow \left(\begin{array}{c} \sigma_1^{i-1} \\ \sigma_2^{i-1} \end{array} \right)$ for $i = 1, \dots, n$.

“Unapplying” a pair of substitutions on a pair of $FOTs$

Rule “**EQUAL FUNCTORS**” uses operation “*unapply*” ‘ \uparrow ’ on a pair of terms t_1, t_2 given a pair of substitutions σ_1, σ_2 :

$$\begin{pmatrix} t_1 \\ t_2 \end{pmatrix} \uparrow \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \stackrel{\text{def}}{=} \begin{cases} \begin{pmatrix} X \\ X \end{pmatrix} & \text{if } t_i = X\sigma_i, \text{ for } i = 1, 2 \\ \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} & \text{otherwise} \end{cases}$$

Declarative \mathcal{FOT} generalization rule for $n = 0$

NB: for $n = 0$, the rule **EQUAL FUNCTORS** becomes an axiom;
viz., for any constant c :

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \vdash \begin{pmatrix} c \\ c \end{pmatrix} c \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}$$

for any pair σ_1, σ_2

\mathcal{FOT} generalization example

Consider the terms $f(a, a, a)$ and $f(b, c, c)$ to generalize; *i.e.*:

- Find term t and substitutions σ_1 and σ_2 such that $t\sigma_1 = f(a, a, a)$ and $t\sigma_2 = f(b, c, c)$:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix} \vdash \begin{pmatrix} f(a, a, a) \\ f(b, c, c) \end{pmatrix} t \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}$$

- By Rule **EQUAL FUNCTORS**, we must have $t = f(u_1, u_2, u_3)$ since:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix} \vdash \begin{pmatrix} f(a, a, a) \\ f(b, c, c) \end{pmatrix} f(u_1, u_2, u_3) \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}$$

where:

- u_1 is the generalization of $\begin{pmatrix} a \\ b \end{pmatrix} \uparrow \begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix}$; that is, of a and b

and by Rule **UNEQUAL FUNCTORS**:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix} \vdash \begin{pmatrix} a \\ b \end{pmatrix} X \begin{pmatrix} \{a/X\} \\ \{b/X\} \end{pmatrix} \text{ therefore } u_1 = X$$

– u_2 is the generalization of $\begin{pmatrix} a \\ c \end{pmatrix} \uparrow \begin{pmatrix} \{a/X\} \\ \{b/X\} \end{pmatrix}$; that is, of a and c ;

and by Rule **UNEQUAL FUNCTORS**:

$$\begin{pmatrix} \{a/X\} \\ \{b/X\} \end{pmatrix} \vdash \begin{pmatrix} a \\ c \end{pmatrix} Y \begin{pmatrix} \{a/X, a/Y\} \\ \{b/X, c/Y\} \end{pmatrix} \text{ therefore } u_2 = Y$$

– u_3 is the generalization of $\begin{pmatrix} a \\ c \end{pmatrix} \uparrow \begin{pmatrix} \{a/X, a/Y\} \\ \{b/X, c/Y\} \end{pmatrix}$; that is, of Y and Y ;

and by Rule **EQUAL VARIABLES**:

$$\begin{pmatrix} \{a/X, a/Y\} \\ \{b/X, c/Y\} \end{pmatrix} \vdash \begin{pmatrix} Y \\ Y \end{pmatrix} Y \begin{pmatrix} \{a/X, a/Y\} \\ \{b/X, c/Y\} \end{pmatrix} \text{ therefore } u_3 = Y$$

• therefore, the overall constraint is thus solved proving the overall judgment valid as:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix} \vdash \begin{pmatrix} f(a, a, a) \\ f(b, c, c) \end{pmatrix} f(X, Y, Y) \begin{pmatrix} \{a/X, a/Y\} \\ \{b/X, c/Y\} \end{pmatrix}$$

i.e., $t = f(X, Y, Y)$, with $\sigma_1 = \{a/X, a/Y\}$ s.t. $t\sigma_1 = f(a, a, a)$,

and $\sigma_2 = \{b/X, c/Y\}$ and $t\sigma_2 = f(b, c, c)$

Going from crisp to fuzzy...

- **extending the foregoing to fuzzy lattice operations as fuzzy constraints**

Fuzzy equivalence relation on a (crisp) set (fuzzy set of pairs)

When S is a finite discrete set $\{x_1, \dots, x_n\}$, since a similarity relation \sim on S is a fuzzy subset of $S \times S$, the three conditions of an equivalence can be visualized on a square $n \times n$ matrix $\sim \subseteq [0, 1]^2$ as follows; $\forall i, j, k = 1, \dots, n$:

- ▶ **reflexivity**: $\sim_{ii} = 1$ entries on the diagonal are equal to 1
- ▶ **symmetry**: $\sim_{ij} = \sim_{ji}$ symmetric entries on either side of the diagonal are equal
- ▶ **transitivity**: $\sim_{ik} \wedge \sim_{kj} \leq \sim_{ij}$ going via an intermediate will always result in a smaller or equal truth value than going directly

N.B.: if $x_i \sim_\alpha x_j$ for some $\alpha \in (0, 1]$, then $x_i \sim_\beta x_j$ for all $\beta \in (0, \alpha]$

Given a similarity relation \sim on signature Σ Sessa extends it homomorphically to *FOTs* as follows:

- ▶ for all $X \in \mathcal{V}$: $X \sim_1 X$
- ▶ for all $X \in \mathcal{V}$ and $t \in \mathcal{T}$ s.t. $t \neq X$: $X \sim_0 t$ and $t \sim_0 X$
- ▶ for $f \in \Sigma_n$ and $g \in \Sigma_n$ s.t. $f \sim_\alpha g$ and $s_i \sim_{\alpha_i} t_i$:

$$f(s_1, \dots, s_n) \sim_{\alpha \wedge \bigwedge_{i=1}^n \alpha_i} g(t_1, \dots, t_n)$$

$$\alpha \in [0, 1], \alpha_i \in [0, 1] \quad (i = 1, \dots, n)$$

Unification degree of pair of terms (0 for dissimilar pairs)

NB: (1) for Sessa's "weak" similarity on Σ : $n \neq m \rightarrow (\sim \cap \Sigma_m \times \Sigma_n = \emptyset)$, for all $m, n \geq 0$
 and (2) operation \wedge is **min** — but other interpretations are possible

Fuzzy subsumption

$$\alpha \in (0, 1]$$

$$t_1 \preceq_{\alpha} t_2$$


iff

$$\exists \sigma : \mathcal{V} \rightarrow \mathcal{T}_{\Sigma, \mathcal{V}}$$

s.t.

$$t_1 \sim_{\alpha} t_2 \sigma$$

Fuzzy unification as a constraint



Fuzzy unification rule

A **fuzzy unification rule** rewrites E_α , a prior set of equations E with truth value $\alpha \in (0, 1]$, into $E'_{\alpha'}$, a posterior set of equations E' with truth value $\alpha' \in [0, \alpha]$, when an optional meta-condition holds:

RULE NAME:

Prior set of equations E_α _____ [Optional meta-condition]

Posterior set of equations $E'_{\alpha'}$

Sessa's "weak" fuzzy unification

VARIABLE ELIMINATION:

$$\frac{(E \cup \{ X \doteq t \})_\alpha}{(E[X \leftarrow t] \cup \{ X \doteq t \})_\alpha} \left[\begin{array}{l} X \notin \mathbf{Var}(t) \\ X \text{ occurs in } E \end{array} \right]$$

CRISP VERSION IS HMM'S:

$$\frac{E \cup \{ X \doteq t \}}{E[X \leftarrow t] \cup \{ X \doteq t \}} \left[\begin{array}{l} X \notin \mathbf{Var}(t) \\ X \text{ occurs in } E \end{array} \right]$$

VARIABLE ERASURE:

$$\frac{(E \cup \{ X \doteq X \})_\alpha}{E_\alpha}$$

CRISP VERSION IS HMM'S:

$$\frac{E \cup \{ X \doteq X \}}{E}$$

EQUATION ORIENTATION:

$$\frac{(E \cup \{ t \doteq X \})_\alpha}{(E \cup \{ X \doteq t \})_\alpha} [t \notin \mathcal{V}]$$

CRISP VERSION IS HMM'S:

$$\frac{E \cup \{ t \doteq X \}}{E \cup \{ X \doteq t \}} [t \notin \mathcal{V}]$$

WEAK TERM DECOMPOSITION:

$$\frac{(E \cup \{ f(s_1, \dots, s_n) \doteq g(t_1, \dots, t_n) \})_\alpha}{(E \cup \{ s_1 \doteq t_1, \dots, s_n \doteq t_n \})_{\alpha \wedge \beta}} \left[\begin{array}{l} f \sim_\beta g \\ n \geq 0 \end{array} \right]$$

NB: only unification rule among HMM's that constrains the overall unification degree upon equating similar terms with different constructors

CRISP VERSION IS ALSO HMM'S:

$$\frac{E \cup \{ f(s_1, \dots, s_n) \doteq f(t_1, \dots, t_n) \}}{E \cup \{ s_1 \doteq t_1, \dots, s_n \doteq t_n \}} [n \geq 0]$$

Fuzzy unification example

Let $\{a, b, c, d\} \subseteq \Sigma_0$, $\{f, g\} \subseteq \Sigma_2$, $\{h\} \subseteq \Sigma_3$; with $a \sim_{.7} b$, $c \sim_{.6} d$, $f \sim_{.9} g$.

- Fuzzy equational constraint to normalize:

$$\{ h(f(a, X_1), g(X_1, b), f(Y_1, Y_1)) \doteq h(X_2, X_2, g(c, d)) \}_1$$

- apply Rule **WEAK TERM DECOMPOSITION** with $\alpha = 1$ and $\beta = 1$:

$$\{ f(a, X_1) \doteq X_2, g(X_1, b) \doteq X_2, f(Y_1, Y_1) \doteq g(c, d) \}_1$$

- apply Rule **EQUATION ORIENTATION** to $f(a, X_1) \doteq X_2$ with $\alpha = 1$:

$$\{ X_2 \doteq f(a, X_1), g(X_1, b) \doteq X_2, f(Y_1, Y_1) \doteq g(c, d) \}_1$$

- apply Rule **VARIABLE ELIMINATION** to $X_2 \doteq f(a, X_1)$ with $\alpha = 1$:

$$\{ X_2 \doteq f(a, X_1), g(X_1, b) \doteq f(a, X_1), f(Y_1, Y_1) \doteq g(c, d) \}_1$$

- apply Rule **WEAK TERM DECOMPOSITION** to $g(X_1, b) \doteq f(a, X_1)$ with $\alpha = 1$ and $\beta = .9$:

$$\{ X_2 \doteq f(a, X_1), X_1 \doteq a, b \doteq X_1, f(Y_1, Y_1) \doteq g(c, d) \}_{.9}$$

- apply Rule **VARIABLE ELIMINATION** to $X_1 \doteq a$ with $\alpha = .9$:
 $\{ X_2 \doteq f(a, a), X_1 \doteq a, b \doteq a, f(Y_1, Y_1) \doteq g(c, d) \}_{.9}$
- apply Rule **WEAK TERM DECOMPOSITION** to $b \doteq a$ with $\alpha = .9$ and $\beta = .7$:
 $\{ X_2 \doteq f(a, a), X_1 \doteq a, f(Y_1, Y_1) \doteq g(c, d) \}_{.7}$
- apply Rule **WEAK TERM DECOMPOSITION** to $f(Y_1, Y_1) \doteq g(c, d)$ with $\alpha = .7$ and $\beta = .9$:
 $\{ X_2 \doteq f(a, a), X_1 \doteq a, Y_1 \doteq c, Y_1 \doteq d \}_{.7}$
- apply Rule **VARIABLE ELIMINATION** to $Y_1 \doteq c$ with $\alpha = .7$:
 $\{ X_2 \doteq f(a, a), X_1 \doteq a, Y_1 \doteq c, c \doteq d \}_{.7}$
- apply Rule **WEAK TERM DECOMPOSITION** to $c \doteq d$ with $\alpha = .7$ and $\beta = .6$:
 $\{ X_2 \doteq f(a, a), X_1 \doteq a, Y_1 \doteq c \}_{.6}$

This is in normal form, yielding substitution σ :

$$\sigma = \{ X_1 = a, Y_1 = c, X_2 = f(a, a) \}$$

with truth value $.6$ so that:

$$t_1\sigma = h(f(a, a), g(a, b), f(c, c)) \sim_{.6} t_2\sigma = h(f(a, a), f(a, a), g(c, d))$$

Moving on to...



fuzzy generalization



Fuzzy generalization judgment

Statement of the form:

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}_\alpha \vdash \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} t \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}_\beta$$

where (for $i = 1, 2$):

- $t \in \mathcal{T}$ and $t_i \in \mathcal{T}$ are \mathcal{FOT} s
- $\sigma_i : \mathcal{V} \rightarrow \mathcal{T}$ are substitutions and $\alpha \in [0, 1]$
- $\theta_i : \mathcal{V} \rightarrow \mathcal{T}$ are substitutions and $\beta \in [0, 1]$

Fuzzy generalization judgment validity

A fuzzy generalization judgment:

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}_\alpha \vdash \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} t \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}_\beta$$


is deemed **valid** whenever ($i = 1, 2$):

$$t_i \sigma_i \sim_\beta t \theta_i$$

with: $0 \leq \beta \leq \alpha \leq 1$

and: $\theta_i \preceq_\beta \sigma_i$ (i.e., $\exists \delta_i$ s.t. $\theta_i \sim_\beta \delta_i \sigma_i$)

Fuzzy generalization judgment validity as a constraint



Fuzzy generalization axioms

FUZZY EQUAL VARIABLES :

$$\left(\begin{array}{c} \sigma_1 \\ \sigma_2 \end{array} \right)_\alpha \vdash \left(\begin{array}{c} X \\ X \end{array} \right) X \left(\begin{array}{c} \sigma_1 \\ \sigma_2 \end{array} \right)_\alpha$$

FUZZY VARIABLE-TERM :

$[t_1 \in \mathcal{V} \text{ or } t_2 \in \mathcal{V}; t_1 \neq t_2; X \text{ is new}]$

$$\left(\begin{array}{c} \sigma_1 \\ \sigma_2 \end{array} \right)_\alpha \vdash \left(\begin{array}{c} t_1 \\ t_2 \end{array} \right) X \left(\begin{array}{c} \sigma_1 \{ t_1 / X \} \\ \sigma_2 \{ t_2 / X \} \end{array} \right)_\alpha$$

DISSIMILAR FUNCTORS :

$[f \not\approx g; m \geq 0, n \geq 0; X \text{ is new}]$

$$\left(\begin{array}{c} \sigma_1 \\ \sigma_2 \end{array} \right)_\alpha \vdash \left(\begin{array}{c} f(s_1, \dots, s_m) \\ g(t_1, \dots, t_n) \end{array} \right) X \left(\begin{array}{c} \sigma_1 \{ f(s_1, \dots, s_m) / X \} \\ \sigma_2 \{ g(t_1, \dots, t_n) / X \} \end{array} \right)_\alpha$$

Fuzzy generalization rule for similar functors

SIMILAR FUNCTORS :

$$[f \sim_{\beta} g; n \geq 0; \alpha_0 \stackrel{\text{def}}{=} \alpha \wedge \beta]$$

$$\frac{\begin{array}{c} \left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right)_{\alpha_0} \vdash \left(\begin{array}{c} s'_1 \\ t'_1 \end{array} \right) u_1 \left(\begin{array}{c} \sigma_1^1 \\ \sigma_2^1 \end{array} \right)_{\alpha_1} \quad \dots \quad \left(\begin{array}{c} \sigma_1^{n-1} \\ \sigma_2^{n-1} \end{array} \right)_{\alpha_{n-1}} \vdash \left(\begin{array}{c} s'_n \\ t'_n \end{array} \right) u_n \left(\begin{array}{c} \sigma_1^n \\ \sigma_2^n \end{array} \right)_{\alpha_n} \end{array}}{\left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right)_{\alpha} \vdash \left(\begin{array}{c} f(s_1, \dots, s_n) \\ g(t_1, \dots, t_n) \end{array} \right) f(u_1, \dots, u_n) \left(\begin{array}{c} \sigma_1^n \\ \sigma_2^n \end{array} \right)_{\alpha_n}}$$

where $\left(\begin{array}{c} s'_i \\ t'_i \end{array} \right) \stackrel{\text{def}}{=} \left(\begin{array}{c} s_i \\ t_i \end{array} \right) \uparrow_{\alpha_i} \left(\begin{array}{c} \sigma_1^{i-1} \\ \sigma_2^{i-1} \end{array} \right)$ for $i = 1, \dots, n$.

Fuzzy “unapplication” of a pair of substitutions on a pair of \mathcal{FOT} s

Rule “**SIMILAR FUNCTORS**” uses operation “**fuzzy unapply**” ‘ \uparrow_α ’ on a pair of terms t_1, t_2 given a pair of substitutions σ_1, σ_2 and truth value $\alpha \in [0, 1]$:

$$\begin{pmatrix} t_1 \\ t_2 \end{pmatrix} \uparrow_\alpha \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \stackrel{\text{def}}{=} \begin{cases} \begin{pmatrix} X \\ X \end{pmatrix} & \text{if } t_i \sim_\alpha X\sigma_i, \text{ for } i = 1, 2 \\ \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} & \text{otherwise} \end{cases}$$

Fuzzy generalization example

Again, let $\{a, b, c, d\} \subseteq \Sigma_0$, $\{f, g\} \subseteq \Sigma_2$, $\{h\} \subseteq \Sigma_3$; with $a \sim_{.7} b$, $c \sim_{.6} d$, $f \sim_{.9} g$.

- Terms to generalize:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix}_1 \vdash \begin{pmatrix} h(f(a, X_1), g(X_1, b), f(Y_1, Y_1)) \\ h(X_2, X_2, g(c, d)) \end{pmatrix} t \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}_\alpha$$

- By Rule **SIMILAR FUNCTORS**, we must have $t = h(u_1, u_2, u_3)$ since:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix}_1 \vdash \begin{pmatrix} h(f(a, X_1), g(X_1, b), f(Y_1, Y_1)) \\ h(X_2, X_2, g(c, d)) \end{pmatrix} h(u_1, u_2, u_3) \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}_\alpha$$

where:

- u_1 is the fuzzy generalization of $\begin{pmatrix} f(a, X_1) \\ X_2 \end{pmatrix} \uparrow_1 \begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix}$; that is, of $f(a, X_1)$ and X_2 ;

by Rule **FUZZY VARIABLE-TERM**:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix}_1 \vdash \begin{pmatrix} f(a, X_1) \\ X_2 \end{pmatrix} X \begin{pmatrix} \{f(a, X_1)/X\} \\ \{X_2/X\} \end{pmatrix}_1 \text{ so } u_1 = X$$

- u_2 is the fuzzy generalization of $\begin{pmatrix} g(X_1, b) \\ X_2 \end{pmatrix} \uparrow_1 \begin{pmatrix} \{f(a, X_1)/X\} \\ \{X_2/X\} \end{pmatrix}$; i.e., $g(X_1, b)$ and X_2 by Rule **FUZZY VARIABLE-TERM**:

$$\begin{pmatrix} \{f(a, X_1)/X\} \\ \{X_2/X\} \end{pmatrix}_1 \vdash \begin{pmatrix} g(X_1, b) \\ X_2 \end{pmatrix} Y \begin{pmatrix} \{\dots, g(X_1, b)/Y\} \\ \{\dots, X_2/Y\} \end{pmatrix}_1 \quad \text{so } u_2 = Y$$

- $u_3 = f(v_1, v_2)$ is the fuzzy generalization of $\begin{pmatrix} f(Y_1, Y_1) \\ g(c, d) \end{pmatrix} \uparrow_{.9} \begin{pmatrix} \{f(a, X_1)/X, g(X_1, b)/Y\} \\ \{X_2/X, X_2/Y\} \end{pmatrix}$; that is, of $f(Y_1, Y_1)$ and $g(c, d)$ with truth value $.9$, because of Rule **SIMILAR FUNCTORS** and $f \sim_{.9} g$, where:

- * v_1 is the fuzzy generalization of $\begin{pmatrix} Y_1 \\ c \end{pmatrix} \uparrow_{.9} \begin{pmatrix} \{f(a, X_1)/X, g(X_1, b)/Y\} \\ \{X_2/X, X_2/Y\} \end{pmatrix}$; i.e., Y_1 and c by Rule **FUZZY VARIABLE-TERM**:

$$\begin{pmatrix} \{f(a, X_1)/X, g(X_1, b)/Y\} \\ \{X_2/X, X_2/Y\} \end{pmatrix}_{.9} \vdash \begin{pmatrix} Y_1 \\ c \end{pmatrix} Z \begin{pmatrix} \{\dots, Y_1/Z\} \\ \{\dots, c/Z\} \end{pmatrix}_{.9} \quad \text{so } v_1 = Z$$

* v_2 is the fuzzy generalization of $\begin{pmatrix} Y_1 \\ d \end{pmatrix} \uparrow_{.9} \begin{pmatrix} \{f(a, X_1)/X, g(X_1, b)/Y, Y_1/Z\} \\ \{X_2/X, X_2/Y, c/Z\} \end{pmatrix}$; i.e., Y_1 and d ; by Rule **FUZZY VARIABLE-TERM**:

$$\begin{pmatrix} \{f(a, X_1)/X, g(X_1, b)/Y, Y_1/Z\} \\ \{X_2/X, X_2/Y, c/Z\} \end{pmatrix} \vdash_{.9} \begin{pmatrix} Y_1 \\ d \end{pmatrix} U \begin{pmatrix} \{\dots, Y_1/U\} \\ \{\dots, d/U\} \end{pmatrix} \text{ so, } v_2 = U$$

in other words, $u_3 = f(Z, U)$ since:

$$\begin{pmatrix} \{f(a, X_1)/X, g(X_1, b)/Y\} \\ \{X_2/X, X_2/Y\} \end{pmatrix} \vdash_1 \begin{pmatrix} f(Y_1, Y_1) \\ g(c, d) \end{pmatrix} f(Z, U) \begin{pmatrix} \{\dots, Y_1/Z, Y_1/U\} \\ \{\dots, c/Z, d/U\} \end{pmatrix}_{.9}$$

Therefore:

$$\begin{pmatrix} \emptyset \\ \emptyset \end{pmatrix} \vdash_1 \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} h(X, Y, f(Z, U)) \begin{pmatrix} \{f(a, X_1)/X, g(X_1, b)/Y, Y_1/Z, Y_1/U\} \\ \{X_2/X, X_2/Y, c/Z, d/U\} \end{pmatrix}_{.9}$$

whereby

$$t\sigma_1 = h(f(a, X_1), g(X_1, b), f(Y_1, Y_1)) = t_1,$$

$$t\sigma_2 = h(X_2, X_2, f(c, d)) \sim_{.9} h(X_2, X_2, g(c, d)) = t_2$$

So we now have fuzzy lattice operations on $\mathcal{FOT}\dots$



but, aren't we missing something?



Hey! ... *but what about similar functors with different arities?*

... *or equal arities but different order of arguments?*

- ▶ **Disallowed in Sessa's weak unification**, even though this would be of great convenience; *e.g.*, in **approximate data retrieval and mining in non-aligned databases**

For example:

person(*Name*, *SSN*, *Address*)

\sim_{α}

individual(*Name*, *DoB*, *SSN*, *Address*)

for $\alpha \in (0, 1]$ would allow fuzzy matching of non-aligned similar records

Similar terms with different argument number or order

Given $\sim : \Sigma^2 \rightarrow [0, 1]$ similarity on $\Sigma \stackrel{\text{def}}{=} \bigcup_{n \geq 0} \Sigma_n$, s.t.:

- $\sim \cap \Sigma_m \times \Sigma_n \neq \emptyset$ for some $m \geq 0, n \geq 0$, with $m \neq n$
- for $f \in \Sigma_m, g \in \Sigma_n, 0 \leq m \leq n$, whenever $f \sim_\alpha g$ there is an *injective mapping* $p : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ that is denoted as $f \sim_\alpha^p g$; e.g.:

$$\begin{array}{c} \textit{person}(\textit{Name}, \textit{SSN}, \textit{Address}) \\ \sim_{.9}^{\{1 \rightarrow 1, 2 \rightarrow 3, 3 \rightarrow 4\}} \\ \textit{individual}(\textit{Name}, \textit{DoB}, \textit{SSN}, \textit{Address}) \end{array}$$

N.B.: m and n are such that $0 \leq m \leq n$; so the one-to-one argument-position mapping goes from the lesser set to the larger set

Unifying similar functors w/ different arg. number/order

GENERIC WEAK TERM DECOMPOSITION :

$$[f \sim_{\beta}^p g; 0 \leq m \leq n]$$

$$(E \cup \{f(s_1, \dots, s_m) \doteq g(t_1, \dots, t_n)\})_{\alpha}$$

$$\left(E \cup \{s_1 \doteq t_{p(1)}, \dots, s_m \doteq t_{p(m)}\} \right)_{\alpha \wedge \beta}$$

FUZZY EQUATION REORIENTATION :

$$[0 \leq n < m]$$

$$(E \cup \{f(s_1, \dots, s_m) \doteq g(t_1, \dots, t_n)\})_{\alpha}$$

$$(E \cup \{g(t_1, \dots, t_n) \doteq f(s_1, \dots, s_m)\})_{\alpha}$$

Generalizing similar functors w/ different arg. number/order

FUNCTOR/ARITY SIMILARITY LEFT :

$$\left[f \sim_{\beta}^p g; 0 \leq m \leq n; \alpha_0 \stackrel{\text{def}}{=} \alpha \wedge \beta \right]$$

$$\frac{\begin{array}{c} \left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right)_{\alpha_0} \vdash \left(\begin{array}{c} s'_1 \\ t'_1 \end{array} \right) u_1 \left(\begin{array}{c} \sigma_1^1 \\ \sigma_2^1 \end{array} \right)_{\alpha_1} \cdots \left(\begin{array}{c} \sigma_1^{m-1} \\ \sigma_2^{m-1} \end{array} \right)_{\alpha_{m-1}} \vdash \left(\begin{array}{c} s'_m \\ t'_m \end{array} \right) u_m \left(\begin{array}{c} \sigma_1^m \\ \sigma_2^m \end{array} \right)_{\alpha_m} \end{array}}{\left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right)_{\alpha} \vdash \left(\begin{array}{c} f(s_1, \dots, s_m) \\ g(t_1, \dots, t_n) \end{array} \right) f(u_1, \dots, u_m) \left(\begin{array}{c} \sigma_1^m \\ \sigma_2^m \end{array} \right)_{\alpha_m}}$$

where $\left(\begin{array}{c} s'_i \\ t'_i \end{array} \right) \stackrel{\text{def}}{=} \left(\begin{array}{c} s_i \\ t_{p(i)} \end{array} \right) \uparrow_{\alpha_i} \left(\begin{array}{c} \sigma_1^i \\ \sigma_2^i \end{array} \right)$ for $i = 1, \dots, m$.

Generalizing similar functors w/ different arg. number/order (ctd.)

FUNCTOR/ARITY SIMILARITY RIGHT :

$$\left[g \sim_{\beta}^p f; 0 \leq n \leq m; \alpha_0 \stackrel{\text{def}}{=} \alpha \wedge \beta \right]$$

$$\frac{\begin{array}{c} \left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right)_{\alpha_0} \vdash \left(\begin{array}{c} s'_1 \\ t'_1 \end{array} \right) u_1 \left(\begin{array}{c} \sigma_1^1 \\ \sigma_2^1 \end{array} \right)_{\alpha_1} \quad \dots \quad \left(\begin{array}{c} \sigma_1^{n-1} \\ \sigma_2^{n-1} \end{array} \right)_{\alpha_{n-1}} \vdash \left(\begin{array}{c} s'_n \\ t'_n \end{array} \right) u_n \left(\begin{array}{c} \sigma_1^n \\ \sigma_2^n \end{array} \right)_{\alpha_n} \end{array}}{\left(\begin{array}{c} \sigma_1^0 \\ \sigma_2^0 \end{array} \right)_{\alpha} \vdash \left(\begin{array}{c} f(s_1, \dots, s_m) \\ g(t_1, \dots, t_n) \end{array} \right) g(u_1, \dots, u_n) \left(\begin{array}{c} \sigma_1^n \\ \sigma_2^n \end{array} \right)_{\alpha_n}}$$

where $\left(\begin{array}{c} s'_i \\ t'_i \end{array} \right) \stackrel{\text{def}}{=} \left(\begin{array}{c} s_{p(i)} \\ t_i \end{array} \right) \uparrow_{\alpha_i} \left(\begin{array}{c} \sigma_1^i \\ \sigma_2^i \end{array} \right)$ for $i = 1, \dots, n$.

OK — we've had enough for now!...



let us recap and conclude



Recapitulation

We **overviewed 3 lattice structures over $FOTs$** (1 crisp and 2 fuzzy), **gave declarative axioms and rules**, and **expressed the 6 corresponding dual lattice operations as constraints**

(✓ indicates original contribution):

▶ **Conventional signature**

• Unification *(Herbrand–Martelli&Montanari’s)*

✓ Generalization *(declarative version of Reynolds–Plotkin’s)*

▶ **Signature with aligned similarity**

• “Weak” fuzzy unification *(Sessa’s)*

✓ “Weak” fuzzy generalization *(dual to Sessa’s)*

▶ **Signature with misaligned similarity**

✓ Full fuzzy unification *(different/mixed arities)*

✓ Full fuzzy generalization *(different/mixed arities)*

Future Work?

► Implement!

- 👉 Java/Scala Libraries
- 👉 Extend Bousi~Prolog?
- 👉 Applications!
- 👉 *Etc., ...*

► OK... But can all this be made more expressive somehow?

Yes! — Extend these results to the lattice of Order-Sorted Feature terms (fuzzy *OSF* constraints?)

We're working on it...



Coming soon to a theater...er **conference** near you!...

Thank You For Your Attention !

Hassan Aït-Kaci



hak@acm.org

Gabriella Pasi



pasi@disco.unimib.it